

Health impact assessment of exposure to PM₁₀ in Ilam city using Air Q software

Shima Tabasideh ¹, Ataallah Vaisi ², Ali Jafari ³, Reza Rezaee ¹, Mahdi Safari ¹, Gilas Hossieni ¹, Roya Ebrahimi ^{1,✉}

1. Department of Environmental Health Engineering, Environmental Health Research Center, Kurdistan University of Medical Sciences, Sanandaj, Iran
2. Ilam Department of Environment, Ilam, Iran
3. Department of Environmental Health Engineering, School of Health and Nutrition, Lorestan University of Medical Sciences, Khorramabad, Iran

Date of submission: 09 Aug 2017, **Date of acceptance:** 05 Dec 2017

ABSTRACT

Air pollution is a major environmental issue in the world that can be harmful for human health. The aim of this study was to determine the health impacts of particulate matter with an aerodynamic diameter $\leq 10\mu\text{m}$ (PM₁₀) in Ilam in 2013 and 2014. The air pollution data were obtained from the Ilam Department of Environmental Protection. The data were analyzed by Excel software, and the annual mortality and morbidity due to cardiovascular and respiratory diseases attributed to PM₁₀ exposure were estimated by Air Q software. According to the results, the average of annual, winter, and summer PM₁₀ concentration was 68.1, 46.7, and 89.3 $\mu\text{g}/\text{m}^3$, respectively, in 2013. In 2014, the average annual, winter, and summer PM₁₀ concentrations were 51.4, 40.3, and 62.6 $\mu\text{g}/\text{m}^3$, respectively. The total mortality, cardiovascular mortality, respiratory mortality, and hospital admissions due to cardiovascular and respiratory diseases in 2013 were estimated at 89, 47, 9, 46, and 119 cases, respectively, and in 2014 they were estimated at 64, 33, 6, 33 and 85 cases, respectively. The results revealed that 9.6% of total mortality in 2013 and 5.8% total mortality in 2014 were associated with concentrations of more than 20 $\mu\text{g}/\text{m}^3$ PM₁₀ concentration. Therefore, it is concluded that exposure to PM₁₀ can increase morbidity and mortality. Thus, urgent and substantial actions to deal with the adverse effects of particulate matters are found to be necessary.

Keywords: air pollution; health impact assessment; PM₁₀; mortality; Air Q software

Introduction

Air pollution is a serious problem around the world that is growing because of industrialization, urbanization, population growth and climate change, especially in developing countries. Particulate matter in the air, as one of the important pollutants and indicators, has attracted a lot of attention.¹⁻³

Several epidemiological studies have demonstrated a relationship between adverse health effects and air pollution and most of these effects include increased mortality and

hospitalization of patients attributed to cardiovascular disease.^{4,5} The World Health Organization (WHO) reported a mortality of 8.4 million annually worldwide and nearly 865,000 premature mortality as health outcomes of air pollution. In addition, according to WHO estimates, nearly 150,000 premature mortality annually in South Asian countries due to cardiovascular and respiratory diseases and lung cancer can be attributed to air pollutants' exposure.^{1,6}

Particulate matter at high concentrations, in addition to health events, results in economic impacts such as severe erosion of buildings and industrial facilities, land and air travel disruption, crops damage and loss, and social consequences such as increased immigration. Particulate matters in the range of 0.005 to 500 μm are transmitted by air, with the main portion

✉ Roya Ebrahimi
ebrahimi83@yahoo.com

Citation: Tabasideh S, Vaisi A, Jafari A, Rezaee A, Safari M, Hossieni G, et al. Health impact assessment of exposure to PM₁₀ in Ilam city using Air Q software. J Adv Environ Health Res 2017; 5(4): 196-204

of them being in the range of 1.0 to 10 μm . Particles with an aerodynamic diameter of 10 μm and smaller are defined as PM_{10} . This PM_{10} fraction, due to the point of entry and settling down in the lower respiratory system, has the most adverse effects on human health.⁷⁻¹⁰

Exposure to high concentrations of PM_{10} can result in both short and long-term adverse health effects.¹¹ These particles can aggravate respiratory diseases such as asthma, emphysema, bronchitis, silicosis, and lung cancer.^{7,12} Relevant management and also public awareness of adverse health effects of air pollution are considered as the first fundamental step in development of strategic programs to control air pollutants. With quantification of air pollution data, the health effects of air pollution are easily estimated and the priority of control measures is defined by comparison with other risk factors. One of the most valid and reliable tools for estimating the short-term effects of air pollutants is the Air Q model introduced by WHO in 2004. In connection with the estimated mortality associated with PM_{10} , several studies have been conducted around the world, including in New Delhi (India), Qin Yang (China), Trieste (Italy) and South Korea. Furthermore, similar studies have been carried out to estimate the health effects of PM_{10} in

cities of Iran including Tehran, Sanandaj, Shiraz, and Tabriz using Air Q models. In the past decade, due to dust storms in the south, south west and west of Iran, these areas have been faced with the phenomenon of the influx of dust and therefore of exposure to high concentrations of PM_{10} .^{1,2,13,14,15}

Ilam city, the center of Ilam province, is located in the west of Iran. In recent years, for various reasons, including geographic location and topography in neighboring Khuzestan Province and bordering with Iraq and also being close to other countries known as sources of the dust (Saudi Arabia, Syria, Kuwait, Yemen), Ilam has been affected by major disasters, air pollution, and dust invasion. Therefore, as a consequence, this study has aims to assess the health outcomes assigned to PM_{10} exposure in Ilam using the Air Q model.

Materials and Methods

Area of study

Ilam, the center of Ilam province, is located in western Iran (latitude 3°63', altitude 46°42') and 1,387 m above sea level (Fig.1). The city has a population of over 213,579 and is fenced with mountains and forested highlands with a mountainous temperate. Ilam is also an important source of oil and gas in the country.¹⁶




Fig. 1. Ilam location in the world and air quality monitoring stations in 2013 and 2014

Air pollution data collection

PM₁₀ data for Ilam was received from the Ilam Department of Environmental Protection between January 1, 2013 and January 1, 2015. Sampling sites based on the location of this department were selected in a way that they covered most areas of the city. After post-processing using Excel software the data were converted to the input file for the Air Q model.

Air Q software

Air Q is a specialized software developed by WHO to assess the health effects of air quality. It can evaluate the potential effects of a criteria pollutant on human health in a given urban area. This model can provide the health effects of pollutants on human health as having such outcomes as mortality, incidence, and prevalence of diseases.^{1,2,13,14,17}

In this model, relative risk, attributable portion, and base incidence are the key factors. The relative risk (RR) represents the ratio of the probability of an event occurring in an exposed population to the probability of the event occurring in a non-exposed population. Furthermore, the term “base incidence” in the model is the number of outcomes based on epidemiological studies conducted by WHO as calculated for a year. The term “attributable portion” is also the percentage of health outcomes directly attributed to the pollution. Attributable portion is calculated using Equation 1 as follows:

$$AP = \frac{\sum\{[RR(c)-1] \times P(c)\}}{\sum[RR(c) \times P(c)]} \quad (1)$$

where AP is the assigned health outcomes, RR (c) represents the health relative risk in Group C, C is the exposed group derived from epidemiological studies, and P (c) is the proportion of Group C or exposed group. The number of cases per population unit (IE) with a base incidence of health consequences (I) is calculated by Equation 2 as follows:

$$IE = I \times AP \quad (2)$$

$$AP = I \times IE \quad (3)$$

Also, having a population size (N), the

estimated number of cases attributable to exposure (NE) can be estimated by Equation 4 as follows:

$$NE = IE \times N \quad (4)$$

In this study, Air Q 2.2.3 software was used with input data of PM₁₀ in the Ilam population two years (January 1, 2013 to January 1, 2015). The output includes indicators such as RR, attributable portion, etc., for all mortality and morality as a result of cardiovascular and respiratory diseases as well as hospitalized patients for cardiovascular and respiratory diseases.¹⁸

Results and Discussion

The results of average and maximum annual and seasonal (summer and winter) PM₁₀ concentrations in 2013 and 2014 for the Keshvari Square Station are presented in Fig. 2. According to the results of this study, the average of annual, winter, and summer concentrations of PM₁₀ in 2013 were 68.115, 46.7 and 89.32 μ/m³, respectively, and in 2014 these values were 51.497, 40.3 and 62.61 μ/m³, respectively. These values are much higher than the annual average standard for PM₁₀ recommended by WHO (20 μ/m³).¹⁹ The maximum annual, winter, and summer concentrations of PM₁₀ in 2013 were 490.63, 165.82 and 490.63 μ/m³, respectively, and in 2014 they were 521.94, 114.36 and 521.94 μ/m³, respectively. As it is evident from the results, the maximum concentration of PM₁₀ occurred in the summer in both the years.

This situation usually occurs in cities in the south and west of Iran and other studies in Iran have shown similar results. As reported by Shahsavani et al., annual average and maximum concentrations of PM₁₀ were 319.6 and 2028 μ/m³ in Ahvaz in April and September, respectively.²⁰ Based on the results of other studies, this situation exists also in the neighboring countries. Draxler et al. reported higher values than the standard in Kuwait, Iraq, and Saudi Arabia. They also reported the highest concentrations PM₁₀ in June.²¹ Accordingly, the increase of PM₁₀ concentrations in summer in Ilam can be assigned to the arid climate of the

Iraq, Kuwait, Saudi Arabia, and Middle East dust storms. These storms are the main

cause of existence of dust in the west and southwest of Iran.²²




Fig. 2. Descriptive statistics of measured particulate matter with aerodynamic diameters $<10 \mu\text{m}$ concentrations in Ilam, Iran, in 2013 (a), 2014 (b)

Fig. 3 presents the per cent of days that people were exposed to different concentrations of PM₁₀. The highest percentage of days that people were exposed to PM₁₀ particles in both 2013 and 2014 were in the range of 40 to 49 $\mu\text{g}/\text{m}^3$. Figure 4 shows the percentage of people who were affected by the short-term effects of various concentrations of PM₁₀ in 2013 and 2014, which was 63% and 78%, respectively, attributed to the particle concentration of less than 99 $\mu\text{g}/\text{m}^3$. The highest number of people affected by short-term effects of various concentrations of PM₁₀ was in the range of 80-89 $\mu\text{g}/\text{m}^3$ in 2013 and in the range of 40-49 $\mu\text{g}/\text{m}^3$

in 2014.

The number of cases attributable to exposure to PM₁₀ of total mortality, mortality from cardiovascular disease, mortality from respiratory diseases, and the number of hospitalized patients of cardiovascular and respiratory diseases are presented in Table 1. The total number of cases of cardiovascular mortality, respiratory mortality and the number of hospitalized patients for cardiovascular disease and respiratory diseases in 2013 as the average RR range, were 9, 46, and 119 cases, respectively, while in 2014, they were 33, 6, 33, and 85 cases, respectively.




Fig. 3. Percentage of days of exposure to different concentrations of particles with an aerodynamic diameter of residents less than 10 microns in Ilam city in 2013 (a) and 2014 (b)

Table 1. Repeated cases (BI), relative risk (RR), the ratio of attributable (AP) estimated annual number of cases Subject to short-term effects of particles with aerodynamic diameter of less than 10 microns

Health endpoints	Year	BI	RR	Attributable proportion in percent (uncertainty range)	Number of cases excess (uncertainty range)
Total mortality	2013	101 3	1.0074 (1.006-1.008)	4.13 (3.4-4.7)	89 (75-102)
	2014	101 3	1.0074 (1.006-1.008)	2.9 (2.4-3.43)	64 (53-73)
Cardiovascular mortality	2013	497	1.008 (1.005-1.018)	4.45 (2.82-9.48)	47 (30-100)
	2014	497	1.008 (1.005-1.018)	3.19 (2.02-6.92)	33 (21-73)
Respiratory mortality	2013	66	1.012 (1.008-1.037)	6.53 (4.45-17.73)	9 (6-24)
	2014	66	1.012 (1.008-1.037)	4.72 (3.19-13.25)	6 (4-18)
Hospital admissions due to cardiovascular diseases	2013	436	1.009 (1.006-1.013)	4.98 (3.37-7.03)	46 (31-65)
	2014	436	1.009 (1.006-1.013)	3.5 (2.4-5.09)	33 (22-47)
Hospital admissions due to respiratory diseases	2013	126 0	1.008 (1.004-1.011)	4.45 (2.71-6.12)	119 (73-164)
	2014	126 0	1.008 (1.004-1.011)	3.19 (1.94-4.42)	85 (52-118)

In Figs. 5 to 9 the cumulative number of cases of mortality, cardiovascular mortality, respiratory mortality, and the annual number of patients hospitalized for cardiovascular disease and respiratory diseases in the range of moderate RR was evaluated for the health effects of PM₁₀.

Based on Figs. 5 to 9, it can be stated that 76% of these cases in 2013 and 83% of cases in 2014 were attributed to PM₁₀ concentrations less than 120 μm^3 . Based on the Air Q model, cumulative number of mortality cases in 2013 and 2014 were 89 and 64, respectively (Fig. 5).




Fig. 4. Percentage of people exposed to different concentrations of particles with an aerodynamic diameter of less than 10 microns in 2013 (a) and 2014 (b)

Now, according to existing reports in Ilam (with a population of over 213, 579 people), the annual death rates in 2013 and 2014 were 982 and 1099 cases, respectively. Based on this model, 5.8% of the mortality cases in 2014 and 9.06% of mortality cases in 2013 are attributable to PM₁₀ particles with higher concentrations than those of the WHO standard. This proportion has also been reported in the provinces neighboring Ilam. Hosseini et al. reported the total number of mortality, cardiovascular mortality, respiratory mortality, and ultimately the number of patients hospitalized for the average relative risk of

cardiovascular disease and respiratory disease as 120, 23, 118, and 305, respectively, in Sanandaj in 2013 where 73% was attributed to PM₁₀ concentrations of less than 129 μm^3 . According to the population of Sanandaj, 11.7% of the mortality was related to PM₁₀ concentrations which were higher than the WHO standard.¹

Similarly, other studies have reported different results elsewhere based on the Air Q model. For instance, Tominz et al. in Trieste, Italy showed that 1.8% of mortality from cardiovascular and 2.5% mortality from respiratory diseases were related to PM₁₀

concentrations that were higher than those of the standard WHO guidelines.²³




Fig. 5. Estimated cumulative number of mortality related to particles with diameter of less than 10 microns in Ilam Air Q model in the years 2013 and 2014




Fig. 6. Estimated cumulative number of mortality caused by cardiovascular diseases related to particles with a diameter less than 10 microns in Ilam using Air Q model in 2013 and 2014




Fig. 7. Estimate the total cumulative number of mortality from respiratory diseases related to particles with a diameter less than 10 microns in Ilam using Air Q model in 2013 and 2014




Fig. 8. Estimates total cumulative number of hospital admissions for respiratory disease related to particles with a diameter less than 10 microns in Ilam using Air Q model in 2013 and 2014




Fig. 9. Estimated total cumulative number of hospital admissions related to cardiovascular patients less than 10 microns in diameter particle model in Ilam using Air Q in 2013 and 2014

Conclusion

The objective of this study was to measure the adverse health effects of PM₁₀ in one of Iran's border provinces with Iraq using Air Q software which shows the effects of pollutants on inhabitants of a certain area. Accordingly, from comparison of results of the present study and results of similar studies, high mortality rate in Ilam during the studied period can be attributed to high concentrations of PM₁₀ and to the number of days with high PM₁₀ concentrations.

Generally speaking, the results of this study indicate that the concentrations of PM₁₀ in Ilam are higher than those of the WHO guidelines. As a conclusion, with regard to the adverse

effects of PM₁₀ on the public health of Ilam and the increase of mortality and cases of hospitalization, we recommend the urgent establishment of necessary policies to reduce exposure to PM₁₀ in the city.

Acknowledgment

This manuscript is extracted from the project approved by the Environmental Health Research Center of Ilam and is funded by the Kurdistan University of Medical Sciences. The authors offer their thanks to the sponsors of the project. The raw data was also obtained from Ilam Department of Environmental Protection, therefore authors greatly acknowledge the department for its contribution.

References

- Hosseini G, Maleki A, Amini H, Mohammadi S, Hassanvand MS, Giahhi O, et al. Health impact assessment of particulate matter in Sanandaj, Kurdistan, Iran. *J Adv Environ Health Res* 2014; 2(1): 54-62.
- Mohammadi MJ, Godini H, Khak MT, Daryanoosh SM, Dobaradaran S, Goudarzi G. An Association between air quality and COPD in Ahvaz, Iran. *Jundishapur J Chronic Dis Care* 2015; 4(1):1-6.
- Zhang LW, Chen X, Xue XD, Sun M, Han B, Li CP, et al. Long-term exposure to high particulate matter pollution and cardiovascular mortality: a 12-year cohort study in four cities in northern China. *Environ Inter* 2014; 62: 41-7.
- Brunekreef B, Holgate ST. Air pollution and health. *The lancet*. 2002;360(9341):1233-42.
- Hosseini G, Teymouri P, Shahmoradi B, Maleki A. Determination of the Concentration and Composition of PM₁₀ during the Middle Eastern Dust Storms in Sanandaj, Iran. *J Res Health Sci* 2015; 15: 182-8.
- López J, Callén M, Murillo R, Garcia T, Navarro M, De la Cruz M, et al. Levels of selected metals in ambient air PM₁₀ in an urban site of Zaragoza (Spain). *Environ Res* 2005; 99(1): 58-67.
- Heidari-Farsani M, Shirmardi M, Goudarzi G, Alavi-Bakhtiarivand N, Ahmadi-Ankali K, Zallaghi E, et al. The evaluation of heavy metals concentration related to PM₁₀ in ambient air of Ahvaz city, Iran. *J Adv Environ Health Res* 2014; 1(2): 120-8.
- Hoseinzadeh E, Samarghandy MR, Ghorbani Shahn F, Roshanaei G, Jafari J. Rate of suspended particulate distribution (PM_{2.5}, PM₁₀ and TSP) in Hamadan main intercity bus stations and its exposure rate. *Health Sys Res* 2012; 8(7): 1245- 54.
- Hosseini G, Teymouri P, Giahhi O, Maleki A. Health Risk Assessment of Heavy Metals in Atmospheric PM₁₀ in Kurdistan University of Medical Sciences Campus. *J Mazandaran Uni Med Sci* 2016; 25(132): 136-46.
- Omidi-Khaniabadi Y, Rashidi R, Goudarzi G, Zare S. Measurement of mass emission values of gaseous pollutants from the stack of Doroud Cement Plant. *J Health Field* 2014; 2(2): 36-42.
- Pisoni E, Volta M. Modeling Pareto efficient PM₁₀ control policies in Northern Italy to reduce health effects. *Atm Environ* 2009; 43(20): 3243-48.
- Reisen VA, Sarnaglia AJ, Reis NC, Lévy-Leduc C, Santos JM. Modeling and forecasting daily average PM 10 concentrations by a seasonal long-memory model with volatility. *Environ Modelling Software* 2014; 51: 286-95.
- Goudarzi G, Angalic KA, Neisia AK, Babaeia AA, Mohammadid B. Estimation of health effects attributed to NO₂ exposure using airq model. *Arc Hygiene Sci* 2012;1(2): 56-66.
- Naddafi K, Hassanvand MS, Yunesian M, Momeniha F, Nabizadeh R, Faridi S, et al. Health impact assessment of air pollution in megacity of Tehran, Iran. *Iran J Environ Health Sci Eng* 2012; 9 (28): 1-7.
- Nourmoradi H, Goudarzi G, Daryanoosh SM, Omidi-Khaniabadi F. Health Impacts of Particulate Matter in Air using AirQ Model in Khorramabad City, Iran. *J Basic Res Med Sci* 2015; 2(2): 44-52.
- Statistical Center of Iran. Estimated population of country cities for. [cited 2011; Available from: URL:<http://www.amar.org.ir/Default.aspx?tabid=1228>.
- Gharehchahi E, Mahvi AH, Amini H, Nabizadeh R, Akhlaghi AA, Shamsipour M, et al. Health impact assessment of air pollution in Shiraz, Iran: a two-part study. *J Environ Health Sci Eng* 2013; 11(11): 1-8.
- Hassanvand MS, Amini H, Yunesian M. Comments on: the evaluation of PM₁₀, PM_{2.5}, And PM₁ concentration FEs during the middle eastern dust (MED) events in Ahvaz, Iran, from April through September 2010. *J Arid Environ* 2013; 97: 1-2.
- Krzyzanowski M. Methods for assessing the extent of exposure and effects of air pollution.

- Occup Environ Med 1997; 54(3): 145-151.
20. Shahsavani A, Naddafi K, Haghhighifard NJ, Mesdaghinia A, Yunesian M, Nabizadeh R, et al. The evaluation of PM 10, PM 2.5, and PM 1 concentrations during the Middle Eastern Dust (MED) events in Ahvaz, Iran, from April through September 2010. *J Arid Environ* 2012; 77: 72-83.
 21. Draxler RR, Gillette DA, Kirkpatrick JS, Heller J. Estimating PM 10 air concentrations from dust storms in Iraq, Kuwait and Saudi Arabia. *Atm Environ* 2001; 35(25): 4315-30.
 22. Goudie A, Middleton NJ. *Desert dust in the global system*: Springer Science & Business Media. Berlin, Heidelberg: Springer; 2006.
 23. Tominz R, Mazzoleni B, Daris F. Estimate of potential health benefits of the reduction of air pollution with PM10 in Trieste, Italy. *Epidemiol Prev* 2005; 29 (3-4): 149-55.